## Table 6-5 Parker Flats Vegetation Cover in Central Coastal Scrub Areas Subject to Vegetation Cutting Conducted in 2008

ESCA RP 2014 Annual Natural Resource Report

Scientific Name			Baseline Data 2008					
	Common Name	Three Transects in Cut Coastal Sage Scrub Vegetation						
		Mean Percent Cover	Standard Deviation	90% Confidence Interval	Mean Relative Cover	Frequency		
Tree Species								
Quercus agrifolia	coast live oak	2.2%	3.1%	5.3%	2.0%	66.7%		
Total Mean Percent Native Tree Cover		2.2%			2.8%			
Shrub and Sub-shrub Specie	es							
Salvia mellifera	black sage	33.0%	11.6%	19.6%	30.3%	100%		
Baccharis pilularis subsp. consagnuinea	coyote brush	14.2%	12.5%	21.1%	13.0%	100%		
Mimulus aurantiacus	sticky monkeyflower	7.5%	3.3%	5.6%	6.8%	100%		
Toxicodendron diversilobum	poison-oak	7.1%	9.1%	15.3%	6.5%	100%		
Artemisia californica	California sagebrush	6.9%	11.9%	20.1%	6.3%	33.3%		
Eriodictyon californicum	California yerba santa	5.1%	3.9%	6.6%	4.6%	100%		
Arctostaphylos tomentosa subsp. tomentosa	shaggy-barked manzanita	1.3%	2.2%	3.7%	1.2%	33.3%		
Ribes speciosum	fuchsia-flowered gooseberry	0.9%	1.5%	2.5%	0.8%	33.3%		
Eriophyllum confertiflorum	golden yarrow	0.0%			0.0%	0%		
Lepechinia calycina	pitcher sage	0.0%			0.0%	0%		
Acmipson glaber	deerweed	0.0%			0.0%	0%		
Total Mean Percent Native Shrub and Subshrub Cover		75.9%			96.9%			
Total Combined Mean Native Cover Between Shrubs and Subshrubs		0.2%	0.2%	0.3%	0.2%	66.7%		
Target Mean Weed Total (Carpobrotus edulis)		0.0%			0.0%	0%		
Total Mean Percent Native Vegetative Cover		78.3%						
Total Mean Percet Bare Ground (Including Masticated Vegetation)		30.8%						
Total Mean Percent Masticated Vegetation (only calculated in 2014)								
Total Mean Percent Bare Ground		30.8%	14.0%	23.7%	28.2%	100%		

**HMP Species in Bold** 

Note: Not all species observed along transects listed in this table

## Table 6-5 Parker Flats Vegetation Cover in Central Coastal Scrub Areas Subject to Vegetation Cutting Conducted in 2008

ESCA RP 2014 Annual Natural Resource Report

Scientific Name	Common Name	Post-activity Data 2012 (Year 3)  Three Transects in Cut Coastal Sage Scrub Vegetation					
		Tree Species	•				
Quercus agrifolia	coast live oak	3.4%	3.1%	5.2%	3.1%	66.7%	
Total Mean Percent Native Tree Cover		3.4%			3.9%		
Shrub and Sub-shrub Specie	es .						
Salvia mellifera	black sage	3.43%	4.1%	7.0%	3.1%	100%	
Baccharis pilularis subsp. consagnuinea	coyote brush	13.61%	7.5%	12.7%	12.4%	100%	
Mimulus aurantiacus	sticky monkeyflower	14.35%	13.2%	22.2%	13.1%	100%	
Toxicodendron diversilobum	poison-oak	3.11%	1.9%	3.3%	2.8%	100%	
Artemisia californica	California sagebrush	0.77%	1.3%	2.2%	0.7%	33.3%	
Eriodictyon californicum	California yerba santa	6.95%	4.5%	7.6%	6.3%	100%	
Arctostaphylos tomentosa subsp. tomentosa	shaggy-barked manzanita	0.00%			0.0%	0.0%	
Ribes speciosum	fuchsia-flowered gooseberry	0.07%	0.1%	0.2%	0.1%	33.3%	
Eriophyllum confertiflorum	golden yarrow	1.10%	1.6%	2.6%	1.0%	67%	
Lepechinia calycina	pitcher sage	0.07%	0.1%	0.2%	0.1%	33%	
Acmipson glaber	deerweed	13.51%	7.7%	13.0%	12.3%	100%	
Total Mean Percent Native Shrub and Subshrub Cover		57.0%			64.9%		
Total Combined Mean Native Cover Between Shrubs and Subshrubs		27.4%	19.6%	33.0%	25.1%	100%	
Target Mean Weed Total (Carpobrotus edulis)		0.0%			0.0%	0%	
Total Mean Percent Native Vegetative Cover		87.8%					
Total Mean Percet Bare Ground (Including Masticated Vegetation)		21.6%					
Total Mean Percent Masticated Vegetation (only calculated in 2014)							
Total Mean Percent Bare Ground		21.6%	8.1%	13.7%	19.8%	100%	

**HMP Species in Bold** Note: Not all species observed alor

## Table 6-5 Parker Flats Vegetation Cover in Central Coastal Scrub Areas Subject to Vegetation Cutting Conducted in 2008

ESCA RP 2014 Annual Natural Resource Report

Scientific Name	Common Name	Post-activity Data 2014 (Year 5)  Three Transects in Cut Coastal Sage Scrub Vegetation					
		Tree Species	•				
Quercus agrifolia	coast live oak	1.9%	1.7%	0.7%	2.8%	66.7%	
Total Mean Percent Native Tree Cover		1.9%			2.7%		
Shrub and Sub-shrub Specie	es						
Salvia mellifera	black sage	4.3%	4.9%	2.1%	6.5%	100%	
Baccharis pilularis subsp. consagnuinea	coyote brush	10.3%	6.9%	2.9%	15.4%	100%	
Mimulus aurantiacus	sticky monkeyflower	11.3%	10.6%	4.5%	16.9%	66.7%	
Toxicodendron diversilobum	poison-oak	3.7%	4.2%	1.8%	5.5%	66.7%	
Artemisia californica	California sagebrush	2.3%	4.0%	1.7%	3.5%	33.3%	
Eriodictyon californicum	California yerba santa	2.9%	2.7%	1.1%	4.3%	66.7%	
Arctostaphylos tomentosa subsp. tomentosa	shaggy-barked manzanita	2.3%	4.0%	1.7%	3.5%	33%	
Ribes speciosum	fuchsia-flowered gooseberry	0.2%	0.3%	0.1%	0.3%	33.3%	
Eriophyllum confertiflorum	golden yarrow	0.3%	0.5%	0.2%	0.4%	33.3%	
Lepechinia calycina	pitcher sage	0.2%	0.3%	0.1%	0.3%	33.3%	
Acmipson glaber	deerweed	0.0%			0.0%	0%	
Total Mean Percent Native Shrub and Subshrub Cover		37.7%			54.6%		
Total Combined Mean Native Cover Between Shrubs and Subshrubs		29.5%	21.4%	9.0%	44.2%	100%	
Target Mean Weed Total (Carpobrotus edulis)		0.0%			0.0%	0%	
Total Mean Percent Native Vegetative Cover		69.1%					
Total Mean Percet Bare Ground (Including Masticated Vegetation)		26.9%					
Total Mean Percent Masticated Vegetation (only calculated in 2014)		19.3%	5.2%	2.2%		100%	
Total Mean Percent Bare Ground		7.6%	7.2%	3.1%		66.7%	

**HMP Species in Bold** Note: Not all species observed alor